



## Short-term effects of high-intensity laser therapy *versus* ultrasound therapy in the treatment of low back pain: a randomized controlled trial

P. FIORE<sup>1</sup>, F. PANZA<sup>2</sup>, G. CASSATELLA<sup>1</sup>, A. RUSSO<sup>1</sup>, V. FRISARDI<sup>3</sup>,  
V. SOLFRIZZI<sup>3</sup>, M. RANIERI<sup>4</sup>, L. DI TEO<sup>5</sup>, A. SANTAMATO<sup>1</sup>

**Background.** Low back pain (LBP) is a common musculoskeletal disorder that is highly prevalent in the general population. Management of this pathology includes numerous interventions depending on pain severity: analgesic, nonsteroidal anti-inflammatory drugs, steroid injections. However, the effect size and duration of symptom relief are limited. Physical therapy (ultrasound [US], laser therapy, manual therapy, interferential current therapy, Back School, aerobic work, therapeutic aquatic exercise acupuncture) have been reported often with mixed results. **Aim.** To evaluate the short-term effectiveness of high-intensity laser therapy (HILT) versus ultrasound (US) therapy in the treatment of LBP. **Design.** Randomized clinical trial. **Setting.** University hospital. **Populations.** Thirty patients with LBP were randomly assigned to a HILT group or a US therapy group.

**Methods.** Study participants received fifteen treatment sessions of HILT or US therapy over a period of three consecutive weeks (five days/week).

**Results.** For the 30 study participants there were no between-group differences at baseline in Visual Analogic Scale (VAS) and Oswestry Low Back Pain Disability Questionnaire (OLBPDQ) scores. At the end of the 3-week intervention, participants in the HILT group showed a significantly greater decrease in pain

<sup>1</sup>Department of Physical Medicine and Rehabilitation University of Foggia, Foggia, Italy

<sup>2</sup>Geriatric Unit and Gerontology-Geriatric Research Laboratory IRCCS Casa Sollievo della Sofferenza San Giovanni Rotondo, Italy

<sup>3</sup>Department of Geriatrics Center for Aging Brain, Memory Unit University of Bari, Bari, Italy

<sup>4</sup>Department of Physical Medicine and Rehabilitation University of Bari, Bari, Italy

<sup>5</sup>OSMAIRM Rehabilitation Center, Laterza, Taranto, Italy

(measured by the VAS) and an improvement of related disability (measured by the OLBPDQ) compared with the group treated with US therapy.

**Conclusion.** Our findings obtained after 15 treatment sessions with the experimental protocol suggested greater effectiveness of HILT than of US therapy in the treatment of LBP, proposing HILT as a promising new therapeutic option into the rehabilitation of LBP.

**KEY WORDS:** Low back pain - Laser therapy - Ultrasonic therapy.

**Fundings.**—None.

**Acknowledgements.**—The present study was already presented at 17<sup>th</sup> European Congress of Physical and Rehabilitation Medicine, Venice, 23-27 May 2010.

**Conflict of interests.**—None.

Received on October 15, 2010.

Accepted for publication on April 14, 2011.

Epub ahead of print on June 8, 2011.

Corresponding author: Dr. A. Santamato, MD, Department of Physical Medicine and Rehabilitation, University of Foggia, Viale Pinto 1, 71100, Foggia, Italy. E-mail: a.santamato@unifg.it